

Advanced Global Atmospheric Gases Experiment (AGAGE)

R.G. PRINN

Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

R.F. WEISS

Scripps Institution of Oceanography, University of California, La Jolla, California 92093

D.M. CUNNOLD

Georgia Institute of Technology, Atlanta, Georgia 30332

P.J. FRASER

CSIRO, Division of Atmospheric Research, Victoria, Australia 3195

P.G. SIMMONDS

University of Bristol, School of Chemistry, Bristol, United Kingdom BS8 ITS

INTRODUCTION

The Advanced Global Atmospheric Gases Experiment (AGAGE) and its predecessors provide continuous high-frequency gas chromatographic measurements of biogenic and anthropogenic gases that are carried out at globally distributed sites in order to quantitatively determine the source and sink strengths and circulation of a large number of chemically and radiatively important long-lived gases. The program that started in 1978 is conveniently divided into three parts associated with three changes in instrumentation: the Atmospheric Lifetime Experiment (ALE), the Global Atmospheric Gases Experiment (GAGE), and AGAGE. Beginning in 1978, these three successive automated, high-frequency, in situ experiments have documented the long-term behavior of the measured concentrations of these important gases over the past 23 years and show both the evolution of latitudinal gradients and the high-frequency variability due to sources and circulation.

AGAGE began over the 1993-1996 time period and continues to the present. It has two instrument components. First, a highly improved gas chromatographic system measures five biogenic/anthropogenic gases (CH_4 , N_2O , CHCl_3 , CO , and H_2) and five anthropogenic gases (CCl_3F , CCl_2F_2 , CH_3CCl_3 , $\text{CCl}_2\text{FCClF}_2$, and CCl_4). Each species is measured 36 times per day with an electron capture detector (ECD), flame ionization detector (FID), and mercuric oxide reduction detector (MRD; this detector is for CO and H_2 and is currently present at only two of the stations, Mace Head, Ireland, and Cape Grim, Tasmania). Second, a gas chromatograph-mass spectrometer (GC-MS) system measures a wide range of hydrochlorofluorocarbons and hydrofluorocarbons: CH_2FCF_3 (HFC-134a), $\text{CH}_3\text{CCl}_2\text{F}$ (HCFC-141b), CH_3CClF_2 (HCFC-142b), etc. These are now serving as interim or permanent alternatives to the chlorofluorocarbons and other long-lived halocarbons regulated by the Montreal Protocol [UNEP, 1996]. Also measured with the GC-MS system are the methyl halides (CH_3Cl , CH_3Br , and CH_3I) and the halons (CBrF_3 and CBrClF_2). AGAGE also includes development and use of new, much more accurate absolute calibrations for most of the measured gases.

The ALE, GAGE, and AGAGE stations have been located in five globally distributed localities: (1) Ireland, first at Adrigole, 52°N, 10°W (1978-1983), then at Mace Head, 53°N, 10°W (1987-present); (2) U.S. West Coast, first at Cape Meares, Oregon, 45°N, 124°W (1979-1989), then at Trinidad Head, California, 41°N, 124°W (1995-present); (3) Ragged Point, Barbados, 13°N, 59°W (1978-present); (4) Cape Matatula, American Samoa, 14°S, 171°W (1978-present); (5) Cape Grim, Tasmania, 41°S, 145°E (1978-present).

Of special significance is the AGAGE operation at Samoa. It enables a direct intercomparison with similar real-time measurements (and also with flask measurements) by the CMDL group. This intercomparison has already aided us in determining the net effect of calibration and instrument differences on the measurements by each group so that the data from both the AGAGE and CMDL networks can be utilized in combination by theoreticians to investigate chemical and meteorological phenomena.

RECENT PROGRESS

The instrumentation and calibrations used in the ALE, GAGE, and AGAGE experiments and a history of the majority of the anthropogenic ozone-depleting and climate-forcing gases in air based on these experiments have been described in detail [Prinn *et al.*, 2000]. We provide estimates of the long-term trends in total chlorine contained in long-lived halocarbons involved in ozone depletion. We summarize interpretations of these measurements using inverse methods to determine trace gas lifetimes and emissions. Finally, we provide a combined observational and modeled reconstruction of the evolution of chlorocarbons by latitude in the atmosphere over the past 60 years, which can be used as boundary conditions for interpreting trapped air in glaciers and oceanic measurements of chlorocarbon tracers of the deep oceanic circulation [Prinn *et al.*, 2000; see also Walker *et al.*, 2000].

A recent analysis of the entire 1978-2000 ALE-GAGE-AGAGE CH_3CCl_3 data set indicates that OH levels in the southern hemisphere are higher than in the northern hemisphere and that OH levels rose between 1978 and 1988,

and then subsequently decreased in 2000 to levels below 1978 values [Prinn *et al.*, 2001]. The 1994-1998 AGAGE CHCl_3 measurements have been analyzed, showing that this gas has a pronounced seasonal cycle (driven largely by OH), a global average baseline mole fraction of 8.9 ppt with no significant trend, and a global lifetime of 6.3 months. Inverse methods indicate emissions are predominantly in the 30°N-90°N region [O'Doherty *et al.*, 2001]. Miller [1998] established the abundance and trends of CHClF_2 and CH_3Br . Huang [1999] produced new estimates of OH using CH_3CCl_3 , CHClF_2 , CH_2FCF_3 , $\text{CH}_3\text{CCl}_2\text{F}$, and CH_3CClF_2 , simultaneously. An analysis of the GAGE-AGAGE measurements of CH_4 for 1985-1999 indicates a decreasing trend, in agreement with previous independent analyses, and has yielded estimates of CH_4 emissions in each semi-hemisphere [Cunnold *et al.*, 2002]. Measurements of CH_3Br from AGAGE and other investigators have been modeled with the 3D Model of Atmospheric Transport and Chemistry (MATCH) [Jensen, 1999]. A comprehensive discussion of the use of Kalman filters in trace gas inverse problems has been published [Prinn, 2000]. A critical analysis of polynomial-based approaches to the CH_3CCl_3 inverse problem has also been published [Prinn and Huang, 2001].

AGAGE GC-MRD measurements of hydrogen at Cape Grim have been reported and interpreted [Simmonds *et al.*, 2000]. GAGE/AGAGE measurements of CCl_3F indicate its global concentrations reached a maximum in 1993 and decayed slightly after that, while CCl_2F_2 levels continued to increase, but only slowly since 1993 [Cunnold *et al.*, 1997; Prinn *et al.*, 2000]. Several AGAGE scientists were lead authors or co-authors in the 1998 ozone assessment [Kurylo *et al.*, 1999; Prinn *et al.*, 1999]. Analysis of regional pollution events in Ireland suggest that industry estimates of the rate of decline in European emissions are too small [Derwent *et al.*, 1998a]. Measurements of CCl_4 from July 1978 to June 1995 show a maximum in its global concentration in 1990, followed by a small decrease since then [Simmonds *et al.*, 1998a]. GC-MS measurements of CH_2FCF_3 , $\text{CH}_3\text{CCl}_2\text{F}$, and CH_3CClF_2 at Mace Head indicate rapid increases in the levels of these CFC replacements, and that industry estimates of HCFC-141b and particularly HCFC-142b emissions appear far too small to explain the observations [Simmonds *et al.*, 1998b]. Measurements of CHClF_2 (HCFC-22) in the Cape Grim archive and at La Jolla have been used to estimate OH concentrations, and this yields values larger than those determined from CH_3CCl_3 (but because of the larger uncertainty in the HCFC-22-based estimate, the difference is not statistically significant) [Miller *et al.*, 1998].

A Lagrangian dispersion model has been used to analyze 1996 Mace Head observations showing that North American sources occasionally produce pollution events at this station, which are about 10% of those due to the European sources [Ryall *et al.*, 1998]. Mace Head data for CH_4 , N_2O , and CO_2 have been analyzed to elucidate European and northern-hemispheric sources [Derwent *et al.*, 1998b]. Wamsley *et al.* [1998] reported measurements of the halon CBrClF_2 . Fraser *et al.* [1999] discussed halon trends and emissions over the past 20 years. Fraser and Prather [1999] reviewed the current prognosis for recovery of the ozone layer. D.S. Cohan, *et al.* (Methyl iodide observations at Cape Grim, Tasmania, using the AGAGE GC-MS instrument, submitted

to *Journal of Atmospheric Chemistry*, 2002) reported AGAGE CH_3I measurements at Cape Grim and interpreted them using trajectory analysis. Dunse *et al.* [2001] reported the results of modeling studies of Cape Grim pollution episodes recorded in AGAGE measurements.

The ALE/GAGE/AGAGE data are available through the Department of Energy-Carbon Dioxide Information Analysis Center (DOE-CDIAC) World Data Center (e-mail to cpd@ornl.gov, Dataset No. DB-1001).

Acknowledgments. In its initial years, the global network (ALE) was supported first by grants from the Chemical Manufacturers Association and later also by the Upper Atmospheric Research Program of the National Aeronautics and Space Administration (NASA). GAGE was subsequently supported by multiple grants from NASA. In its latest phase (AGAGE) support came (and comes) primarily from NASA (grants NAGW-732, NAG1-1805, NAG5-3974 to the Massachusetts Institute of Technology; grants NAGW-2034, NAG5-4023 to Scripps Institution of Oceanography), with important contributions also from the United Kingdom Department of Environment (now Department of Environment, Transport, and the Regions) (contracts PECD 7/10/154 and EPG 1/1/37 and 1/1/82 to Insccon), Commonwealth Scientific and Industrial Research Organization (Australia), Bureau of Meteorology (Australia), and the Alternative Fluorocarbons Environmental Acceptability Study. Support for the Barbados Station during GAGE/AGAGE has been shared approximately equally between NASA and NOAA. We also thank NOAA for its infrastructure support at the Samoa station by Cooperative Agreement.

REFERENCES

- Cunnold, D.M., R.F. Weiss, R.G. Prinn, D.E. Hartley, P.G. Simmonds, P.J. Fraser, B.R. Miller, F.N. Alyea, and L. Porter, GAGE/AGAGE measurements indicating reductions in global emissions of CCl_3F and CCl_2F_2 in 1992-1994, *J. Geophys. Res.*, **102**, 1259-1269, 1997.
- Cunnold, D.M., *et al.*, GAGE/AGAGE measurements of methane at five sites from 1985-1999 and source inferences there from, *J. Geophys. Res.*, in press, 2002.
- Derwent, R., P.G. Simmonds, S. O'Doherty, and D. Ryall, The impact of the Montreal Protocol on halocarbon concentrations in northern hemisphere baseline and European air masses at Mace Head, Ireland, from 1987-1996, *Atmos. Environ.*, **32**, 3689-3702, 1998a.
- Derwent, R., P.G. Simmonds, S. O'Doherty, P. Ciais, and D. Ryall, European source strengths and Northern Hemisphere baseline concentrations of radiatively active trace gases at Mace Head, Ireland, *Atmos. Environ.*, **32**, 3703-3715, 1998b.
- Dunse, B.L., L.P. Steele, P.J. Fraser, and S.R. Wilson, An analysis of Melbourne pollution episodes observed at Cape Grim from 1995-1998, in *Baseline 97-98*, pp. 34-42, CSIRO-DAR and Bureau of Meteorology, Aspendale, Australia, 2001.
- Fraser, P.J., and M.J. Prather, An uncertain road: The science of atmospheric recovery, *Nature*, **398**, 663-664, 1999.
- Fraser, P.J., D. Oran, C. Reeves, S. Penkett, and A. McCulloch, Southern hemispheric halon trends (1978-1998) and global halon emissions, *J. Geophys. Res.*, **104**, 15,985-15,999, 1999.
- Huang, J., Optimal determination of global tropospheric OH concentrations using multiple trace gases, Ph.D. dissertation, 163 pp., Mass. Inst. of Technol., Cambridge, 1999.
- Jensen, C., Terrestrial sources and sinks of atmospheric methyl bromide: Three-dimensional modeling of tropospheric abundance and sensitivities, M.S. dissertation, *MIT Center for Global Change Science Report* 62, 41 pp., Mass. Inst. of Technol., Cambridge, 1999.
- Kurylo, M.J., *et al.*, Short-lived ozone-related compounds, in *Scientific Assessment of Ozone Depletion: 1998, Global Ozone Res. and Monit. Proj. Rep.* 44, pp. 2.1-2.56, World Meteorol. Org., Geneva, 1999.

- Miller, B.R., Abundances and trends of atmospheric chlorodifluoromethane and bromomethane, Ph.D. dissertation, 149 pp., Univ. of Calif., San Diego, 1998.
- Miller, B.R., J. Huang, R.F. Weiss, R.G. Prinn, and P.J. Fraser, Atmospheric trend and lifetime of chlorodifluoromethane (HCFC-22) and the global tropospheric OH concentration, *J. Geophys. Res.*, **103**, 13,237-13,248, 1998.
- O'Doherty, S., et al., In situ chloroform measurements at AGAGE atmospheric research stations from 1994-1998, *J. Geophys. Res.*, **106**, 20,429-20,444, 2001.
- Prinn, R.G., Measurement equation for trace chemicals in fluids and solution of its inverse, in *Inverse Methods in Global Biogeochemical Cycles*, edited by P. Kasibhatla et al., pp. 3-18, *Geophysical Monograph, No. 114*, Am. Geophys. Union, Washington, DC, 2000.
- Prinn, R.G., and J. Huang, Comment on "Global OH trend inferred from methylchloroform measurements" by M. Krol et al., *J. Geophys. Res.*, **106**, 23,151-23,158, 2001.
- Prinn, R.G., R. Zander, D.M. Cunnold, J.W. Elkins, A. Engel, P.J. Fraser, M.R. Gunson, M.K.W. Ko, E. Mahieu, P.M. Midgley, J.M. Russell III, C.M. Volk, and R.F. Weiss, Long-lived ozone-related compounds, in *Scientific Assessment of Ozone Depletion: 1998, Global Ozone Res. and Monit. Proj. Rep. 44*, pp. 1.1-1.54, World Meteorol. Org., Geneva, 1999.
- Prinn, R.G., R.F. Weiss, P.J. Fraser, P.G. Simmonds, D.M. Cunnold, F.N. Alyea, S. O'Doherty, P. Salameh, B.R. Miller, J. Huang, R.H.J. Wang, D.E. Hartley, C. Harth, L.P. Steele, G. Sturrock, P.M. Midgley, and A. McCulloch, A history of chemically and radiatively important gases in air deduced from ALE/GAGE/AGAGE, *J. Geophys. Res.*, **115**, 17,751-17,792, 2000.
- Prinn, R.G., J. Huang, R. Weiss, D. Cunnold, P. Fraser, P. Simmonds, A. McCulloch, P. Salameh, S. O'Doherty, R. Wang, C. Harth, L. Porter, and B. Miller, Evidence for significant variations in atmospheric hydroxyl radicals over the last two decades, *Science*, **293**, 1048-1054, 2001.
- Ryall, D., R. Maryon, R. Derwent, and P.G. Simmonds, Modeling long-range transport of CFCs to Mace Head, Ireland, *Q. J. R. Meteorol. Soc.*, **124**, 417-446, 1998.
- Simmonds, P.G., D.M. Cunnold, R.F. Weiss, R.G. Prinn, P.J. Fraser, and A. McCulloch, Global trends and emission estimates of carbon tetrachloride (CCl₄) from in situ background observations from July 1978 to June 1995, *J. Geophys. Res.*, **103**, 16,017-16,027, 1998a.
- Simmonds, P.G., J. Huang, R. Prinn, S. O'Doherty, R. Derwent, D. Ryall, G. Nickless, and D. Cunnold, Calculated trends and atmospheric abundance of 1,1,1,2-tetrafluoroethane, 1,1-dichloro-1-fluoroethane, and 1-chloro-1,1-fluoroethane using automated in situ gas chromatography-mass spectrometry measurements recorded at Mace Head, Ireland, from October 1994 to March 1997, *J. Geophys. Res.*, **103**, 16,029-16,037, 1998b.
- Simmonds, P.G., R.G. Derwent, S. O'Doherty, D.B. Ryall, L.P. Steele, R.J. Langenfelds, P. Salameh, H.J. Wang, C.H. Dimmer, and L.E. Hudson, Continuous high frequency observations of hydrogen at the Mace Head baseline atmospheric monitoring station over the 1994-1998 period, *J. Geophys. Res.*, **105**, 12,105-12,121, 2000.
- UNEP (United Nations Environmental Programme), *Handbook for the International Treaties for the Protection of the Ozone Layer*, 4th ed., Nairobi, Kenya, 1996.
- Walker, S.J., R.F. Weiss, and P.K. Salameh, Reconstructed histories of the annual mean atmospheric mole fractions for the halocarbons CFC-11, CFC-12, CFC-113, and carbon tetrachloride, *J. Geophys. Res.*, **105**, 14,285-14,296, 2000.
- Wamsley, P., et al., Distribution of halon-1211 (CBrClF₂) in the upper troposphere and lower stratosphere and the 1994 bromine budget, *J. Geophys. Res.*, **103**, 1513-1526, 1998.